


09/7/9, 272



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

B

Search

PubMed

for

Limits

Preview/Index

History

Clipboard

Details

Preview

Go

C

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

- Search History will be lost after one hour of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.

Search

Most Recent Queries

Time Result

#3 Related Articles for PubMed (Select 7769120)

09:03:39

181

#1 Search j clin invest[jour] AND 95[volume] AND 2806[page] Field: Title Word


08:47:51

1

Clear History

Write to the Help DeskNCBI | NLM | NIHDepartment of Health & Human ServicesFreedom of Information Act | Disclaimer

Apr 24 2003 12:56:45



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

B

Search

PubMed

for

Go

Clear

About Entrez

Limits

Preview/Index

History

Clipboard

Details

Text Version

Display

Abstract

Show:

20

Sort

Send to

Text

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: J Clin Invest 1995 Jun;95(6):2806-12

Related Articles, Li

Increased abundance of the receptor-type protein-tyrosine phosphatase LAR accounts for the elevated insulin receptor dephosphorylating activity in adipose tissue of obese human subjects.

Ahmad F, Considine RV, Goldstein BJ.

Dorrance H. Hamilton Research Laboratories, Department of Medicine, Jefferson Medical College, Thomas Jefferson University, Philadelphia, Pennsylvania 19107, USA.

Protein-tyrosine phosphatases (PTPases) have an essential role in the regulation of the steady-state phosphorylation of the insulin receptor and other proteins in the insulin signalling pathway. To examine whether increased PTPase activity is associated with adipose tissue insulin resistance in human obesity we measured PTPase enzyme activity towards the insulin receptor in homogenates of subcutaneous adipose tissue from a series of six lean and six nondiabetic, obese (body mass index > 30) subjects. The obese subjects had a mean 1.74-fold increase in PTPase activity ($P < 0.0001$) with a striking positive correlation by linear regression analysis between PTPase activity and body mass index among all of the samples ($R = 0.918$; $P < 0.0001$). The abundance of three candidate insulin receptor PTPases in adipose tissue was also estimated by immunoblot analysis. The most prominent increase was a 2.03-fold rise in the transmembrane PTPase LAR ($P < 0.001$). Of the three PTPases examined, only immunodepletion of LAR protein from the homogenates with neutralizing antibodies resulted in normalization of the PTPase activity towards the insulin receptor demonstrating that the increase in LAR was responsible for the enhanced PTPase activity in the adipose tissue from obese subjects. These studies suggest that increased PTPase activity towards the insulin receptor is a pathogenetic factor in the insulin resistance of adipose tissue in human obesity and provide